



## Problem Set 6

Ch153a – Winter 2024

Due: 12 February 2024

- (15 points) Consider a binuclear metal complex constructed from two square-planar  $ML_4$  fragments where the L ligands are  $\sigma$ -donors. There are two limiting conformations in the resulting  $M_2L_8$  complex: in one the ligands are eclipsed and in the other they are staggered.
  - Draw the two conformations of the binuclear metal complex and assign each to a symmetry point group.
  - Construct an MO diagram for each conformation using the following orbitals: five M  $d$  orbitals and four L  $\sigma$  orbitals. Assume that M-L  $\sigma$ -bonding is quite strong.
  - Assume that  $ML_4$  has a  $d^4$  electron configuration, and that each ligand L contributes two  $\sigma$  electrons.
    - Predict the preferred ground-state conformation of the corresponding  $M_2L_8$  complex and give the electronic configuration and term symbol for the ground electronic state. Determine the metal-metal bond order.
    - Identify the spin-allowed electronic transitions involving the  $d$ -orbitals, determine the term symbols for the excited states, and predict the relative energy ordering of these states.
    - Predict the lowest-energy electric-dipole-allowed transition. Also, predict the polarization of light that will induce the transition.
  - Assume that  $ML_4$  has a  $d^6$  electron configuration, and that each ligand L contributes two  $\sigma$  electrons. Predict the preferred ground-state conformation of the corresponding  $M_2L_8$  complex and give the electronic configuration and term symbol for the ground electronic state. Determine the metal-metal bond order.
  - Assume that  $ML_4$  has a  $d^7$  electron configuration, and that each ligand L contributes two  $\sigma$  electrons.
    - Predict the preferred ground-state conformation of the corresponding  $M_2L_8$  complex and give the electronic configuration and term symbol for the ground electronic state. Determine the metal-metal bond order.
    - Identify the spin-allowed electronic transitions involving the  $d$ -orbitals, determine the term symbols for the excited states, and predict the relative energy ordering of these states.
    - Predict the lowest-energy electric-dipole-allowed transition that should be M—M axis polarized.






**Figure 2.** Electronic spectra of  $\text{Mn}_2(\text{CO})_{10}$  in 3-PIP: —, 300 K; - - -, 77 K.


- (5 points) The absorption spectrum of  $\text{Mn}_2(\text{CO})_{10}$  (shown above) exhibits an intense absorption band at  $30,000 \text{ cm}^{-1}$ , and a lower-energy feature at  $27,000 \text{ cm}^{-1}$ . The  $30,000\text{-cm}^{-1}$  absorption band is polarized parallel to the Mn-Mn axis, and the  $27,000 \text{ cm}^{-1}$  band is polarized perpendicular to this axis. Propose assignments for these bands based on a MO diagram for binuclear  $d^7\text{-}d^7$  complexes.
- (10 points) The room-temperature absorption spectrum of a single crystal of  $\text{K}_4\text{Pt}_2(\text{H}_2\text{P}_2\text{O}_5)_4 \cdot 2\text{H}_2\text{O}$  ( $\text{K}_4\text{Pt-POP}$ ) is shown to the right.  $\text{Pt-POP}^{4-}$  also is luminescent, displaying intense green phosphorescence with a maximum near 500 nm.

At cryogenic temperatures, the absorption spectrum of the  $\text{Ba}^{2+}$  salt of  $\text{Pt-POP}^{4-}$  develops rich vibrational fine structure in the lowest energy absorption band (next page). The vibrational fine structure is associated with an excited-state distortion along the Pt-Pt stretching coordinate. The vibronic peak positions for the lowest energy absorption band are given in the table on the following page.



**Figure 2.** Absorption spectrum of a single crystal of  $\text{K}_4\text{Pt}_2(\text{H}_2\text{P}_2\text{O}_5)_4 \cdot 2\text{H}_2\text{O}$  at 300 K. Solid line: electric dipole perpendicular to the crystallographic  $c$  axis. Dashed line: parallel to the  $c$  axis and the Pt-Pt axis.

**Figure 1.** Absorption spectrum along two extinction directions of the same face of a single crystal of  $\text{Ba}_2\text{Pt}_2(\text{H}_2\text{P}_2\text{O}_5)_4$  at 5 K. The orientation of the metal-metal direction is not known in this salt.




- What is the vibrational frequency of the distorting mode in the excited state?
- What value of the Huang-Rhys parameter ( $S_{\text{HR}}$ ) gives the best fit to the lowest energy absorption band?
- If the force constant for the Pt-Pt stretching mode is  $1.0 \text{ mdyne}/\text{\AA}$ , what is the magnitude of the distortion along the Pt-Pt coordinate in the excited state?

**Ba<sub>2</sub>Pt-POP Absorption**

v	$\lambda$ , nm
0	476.0
1	472.5
2	469.1
3	465.7
4	462.4
5	459.2
6	456.1
7	452.9
8	449.8
9	446.7
10	443.5
11	440.4
12	437.4
13	434.4
14	431.5
15	428.7
16	425.8
17	423.2
18	420.3
19	417.5

Figure 5.  ${}^3A_{2u} \rightarrow {}^1A_{1g}$  (phosphorescence) spectrum of  $Ba_2Pt_2(H_2P_2O_5)_4$  at 10 K (bottom).



The phosphorescence spectrum of crystalline  $Ba_2Pt$ -POP at 10 K also displays rich fine structure in the Pt-Pt vibrational mode (below). The vibronic peak positions for the phosphorescence band are given in the table on the following page.

- What is the vibrational frequency of the distorting mode in the ground state?
- What value of the Huang-Rhys parameter ( $S_{HR}$ ) gives the best fit to the phosphorescence band?
- On the basis of your fit to the phosphorescence spectrum, what is the magnitude of the distortion along the Pt-Pt coordinate in the excited state? How does this value compare to that extracted from the fit to the absorption spectrum?
- The Pt-Pt distance in the ground state of  $Pt-POP^{4-}$  is 2.92 Å. On the basis of the structured absorption and phosphorescence band profiles, what do you estimate for the Pt-Pt distance in the excited state?

**Ba<sub>2</sub>Pt-POP  
Phosphorescence**

v	$\lambda$ , nm
0	476.5
1	479.0
2	481.6
3	484.2
4	486.8
5	489.5
6	492.3
7	495.1
8	497.8
9	500.6
10	503.3
11	506.1
12	509.1
13	512.0
14	515.0
15	517.9
16	520.9
17	523.9
18	526.9
19	530.1
20	533.3
21	536.5